



USP

Some more mathematical formulation of stock dynamics

■ Purpose of slides

- Introduce the basics in mathematical representation of population dynamics in some detail
 - Stock production models
 - A revision
 - Cohort based models (generic length/age models)
 - An introduction
- Show how the models are all a special form of the mass balance equation (Russels equation)

■ Source:

- Haddon 2001: Chapter 1 & 2
- Hilborn and Walters 1992: Chapter 3.4

Russel's mass balance formulation

$$\left(\begin{array}{c} \text{next} \\ \text{biomass} \end{array} \right) = \left(\begin{array}{c} \text{last} \\ \text{biomass} \end{array} \right) + \text{recruitment} + \text{growth} - \left(\begin{array}{c} \text{natural} \\ \text{mortality} \end{array} \right) - \text{catch}$$


■ Russels contribution:

- “.. the sole value of the exact formulation given above is that it distinguishes the separate factors making up gain and loss respectively, and is therefore **an aid to clear thinking**” (Russel 1931)
- Recognized that a stock could be divided into animals that were in the fishable stock and those that were entering the fishable stock at any one time (recruitment)
- Stock biomass has **gains**: Recruitment and growth
- Stock biomass has **losses**: Natural and fishing mortality (catch)

Russel's equation: A mass balance equation

- $B_{t+1} = B_t + R_t + G_t - M_t - Y_t$

- B_{t+1} stock size in weight at start of time $t+1$
- B_t stock size in weight at start of time t
- R_t weight of all recruits entering stock at time t
 - Recruits: Young fish "entering" the stock in each time period
- G_t weight increase of fish surviving from t to $t+1$
- M_t weight loss of fish that died from t to $t+1$
- Y_t weight of fish captured from t to $t+1$



USP

Stock production models

Russel equation modified

- Russel equation for biomass:

$$B_{t+1} = B_t + R_t + G_t - M_t - Y_t$$

- In the absence of fishing:

$$B_{t+1} = B_t + R_t + G_t - M_t$$

- The two sources of **increase** are called **production**:

$$\text{Production} = (R_t + G_t)$$

- Difference between production and natural mortality is called **surplus production**:



$$\text{Surplus production} = (R_t + G_t) - M_t$$

- If the processes of recruitment, growth and natural mortality are constant we can write the Russel equation as:

$$B_{t+1} = B_t + rB_t$$

- r : intrinsic growth rate
 - Here recruitment, growth and mortality are all lumped into one number
 - $r > 0$, population will grow
 - $r = 0$, population remains constant
 - $r < 0$, population decreases with time



Population growth curves



■ Model limitations

- No population increase or decreases continuously and there seems to be an upper bound due to food / space limitation, predation, competition. This model is thus not realistic to describe **long term** population change
- Note this is an exponential model

Density dependent model



- Alter the exponential model by taking into account that population growth rates is a function of population size:
 - $r' = r_0 - r_1B$
 - where r_0 is the population growth rate when the population size is small (mathematically NULL)
 - r_1 is a value that scale the rates with population size (B)

By mathematical derivation, taking into account a linear density dependent effect of birth and death rate, we can expand the exponential model:


$$B_{t+1} = B_t + rB_t$$

to the following form

$$B_{t+1} = B_t + r \left(\frac{K - B_t}{K} \right) B_t$$

- K : The carrying capacity (often written as B_{\max}).
- r : The intrinsic rate of growth (r):
 - is multiplied by the difference between the current population size and the carrying capacity ($K - B_t / K$).

Population trajectory according to logistic model



$$B_{t+1} = B_t + r \left(\frac{K - B_t}{K} \right) B_t$$

Time t

Why do we want to know about r and K ?

		r		K				
		0.25		1000 tonnes				
t	Bt (tonnes)	Bt/K	1-Bt/K	r	$r(1-Bt/K)$	Surplus production (tonnes)		
1	15	0.02	0.99	0.25	0.25	4		
2	19	0.02	0.98	0.25	0.25	5		
3	23	0.02	0.98	0.25	0.24	6		
4	29	0.03	0.97	0.25	0.24	7		
5	36	0.04	0.96	0.25	0.24	9		
...								
18	431	0.43	0.57	0.25	0.14	61		
19	492	0.49	0.51	0.25	0.13	62		
20	554	0.55	0.45	0.25	0.11	62		
21	616	0.62	0.38	0.25	0.10	59		
...								
41	998	1.00	0.00	0.25	0.00	1		
42	998	1.00	0.00	0.25	0.00	0		
43	999	1.00	0.00	0.25	0.00	0		


When $Bt \ll K$, the term in the brackets $\rightarrow 1$
 When $Bt \rightarrow K$, the term in the bracket $\rightarrow 0$

Higher r , faster population growth and Surp. Pr.

t	r		K		Surplus production (tonnes)	
	Bt (tonnes)	Bt/K	1-Bt/K	r	r(1-Bt/K)	
		1.00				
		1000 tonnes				
1	15	0.02	0.99	1.00	0.99	15
2	30	0.03	0.97	1.00	0.97	29
3	59	0.06	0.94	1.00	0.94	55
4	114	0.11	0.89	1.00	0.89	101
5	215	0.21	0.79	1.00	0.79	169
6	383	0.38	0.62	1.00	0.62	236
7	620	0.62	0.38	1.00	0.38	236
8	856	0.86	0.14	1.00	0.14	124
9	979	0.98	0.02	1.00	0.02	20
10	1000	1.00	0.00	1.00	0.00	0

For a given K , the higher the growth rate the faster the population response and the greater the MSY

Production as a function of stock size



Are related directly to Russels mass-balance formulation:

$$B_{t+1} = B_t + R_t + G_t - M_t - Y_t$$

$$B_{t+1} = B_t + P_t - Y_t$$

$$B_{t+1} = B_t + f(B_t) - Y_t$$

- B_{t+1} : Biomass in the beginning of year $t+1$ (or end of t)
- B_t : Biomass in the beginning of year t
- P_t : Surplus production
 - the difference between production (recruitment + growth) and natural mortality
- $f(B_t)$: Surplus production as a function of biomass in the start of the year t
- Y_t : Biomass (yield) caught during year t

- Classic Schaefer (logistic) form:


$$f B_t = rB_t \left(1 - \frac{B_t}{K} \right)$$

- The more general Pella & Tomlinson form:

$$f B_t = \frac{r}{p} B_t \left(1 - \left[\frac{B_t}{K} \right]^p \right)$$

- Note: when $p=1$ the two functional forms are the same
- If $p \neq 1$, then the density dependence is no longer linear


Pella-Tomlinson model: influence of p value



$$B_{t+1} = B_t + r \left(\frac{K - B_t}{K} \right) B_t - Y_t$$

$$CPUE_t = qB_t$$

- Need a time series of:
 - Total annual catch (Y_t)
 - Index of abundance ($CPUE_t$)



USP

Cohort models

Russel equation modified 1

- Russel equation for biomass:

$$B_{t+1} = B_t + R_t + G_t - M_t - Y_t$$

- Russel equation for numbers:

$$N_{t+1} = N_t + R_t - D_t - C_t$$

- N_{t+1} stock size in numbers at start of time $t+1$
 - N_t stock size in numbers at start of time t
 - R_t **number** of recruits entering the stock at time t
 - D_t number of fish that died from t to $t+1$
 - C_t number of fish that are caught from t to $t+1$
- What happened to the growth term??

Russel equation modified 2

- Russel equation for numbers:

$$N_{t+1} = N_t + R_t - D_t - C_t$$

- If we are considering ONE cohort only we can drop the recruitment term and have

$$N_{t+1} = N_t - D_t - C_t$$

$$\left(\begin{array}{c} \text{Numbers alive} \\ \text{at the beginning} \\ \text{of next time period} \end{array} \right) = \left(\begin{array}{c} \text{Number alive} \\ \text{at the beginning} \\ \text{of this time period} \end{array} \right) - \left(\begin{array}{c} \text{Numbers dying} \\ \text{naturally} \\ \text{this period} \end{array} \right) - \left(\begin{array}{c} \text{Catch} \\ \text{this} \\ \text{period} \end{array} \right)$$

- If we assume that the total number dying ($D_t + C_t$) and are a proportion of those living we have

$$N_{t+1} = N_t - mN_t = N_t (1 - m) = sN_t$$

- m : proportion of fish that die during time interval t to $t+1$
- s : proportion of fish that survive during time interval t to $t+1$
- $s+m = 1$

- The equation:

$$N_{t+1} = N_t (1 - m) = sN_t$$

is an exponential model and the discrete version is:


$$N_{t+1} = N_t e^{-Z_t}$$

i.e. the negative exponential model

m & s: proportional coefficients

Z: instantaneous coefficient


After birth there is only death!



- After hatching the individuals in a cohort can only decline
- If this does not hold true the stock is wrongly defined

$$N_t = N_0 e^{-Zt}$$

The higher the Z the faster
the population numbers decline



- Since mortality is not constant through out a cohorts life we work in smaller time steps:

$$N_{t+1} = N_t e^{-Z_t}$$

- N_t : Number of fish at age time t
- N_{t+1} : Number of fish at time $t+1$
- Z_t : instantaneous mortality coefficient over time period t to $t+1$

Separation of fishing and natural mortality


- Since we are often interested in separating natural and fishing mortality we write:

$$N_{t+1} = N_t e^{-M_t + F_t}$$

- M_t : natural mortality of at time t
- F_t : fishing mortality of at time t
- We will refer to this equation as the stock equation.

The effect of fishing

$$N_{t+1} = N_t e^{-M_t + F_t}$$

M=0.2

Describing the catch 1

- The number of fish that die in each time interval is:

$$D_t = N_t - N_{t+1}$$

- Substituting with the stock equation we get:

$$\begin{aligned} D_t &= N_t - N_t e^{-M_t + F_t} \\ &= N_t (1 - e^{-M_t + F_t}) \end{aligned}$$

$$\left(\begin{array}{l} \text{Numer of} \\ \text{fish that die} \\ \text{during the time} \end{array} \right) = \left(\begin{array}{l} \text{Number of} \\ \text{fish alive in} \\ \text{the beginning} \end{array} \right) \left(\begin{array}{l} \text{Proportion of} \\ \text{fish that die} \end{array} \right)$$

- The number that die due to fishing mortality is the fraction (F_t/Z_t) of the number of fish that die, i.e.

$$C_t = \frac{F_t}{F_t + M_t} (1 - e^{-F_t + M_t}) N_t$$

$$\left(\begin{array}{l} \text{Numer of} \\ \text{fish fished} \\ \text{during the time} \end{array} \right) = \left(\begin{array}{l} \text{Proportion of} \\ \text{fish that die} \\ \text{due to fishing} \end{array} \right) \left(\begin{array}{l} \text{Proportion of} \\ \text{fish that die} \end{array} \right) \left(\begin{array}{l} \text{Number of} \\ \text{fish alive in} \\ \text{the beginning} \end{array} \right)$$

- C_t : The number of fish caught over time t to $t+1$

Describing the catch 3

- It can be shown that if we take the average population size of the period t to $t+1$ (\bar{N}_t) the catch equation becomes:

$$C_t = F_t \bar{N}_t$$

- The form of the stock equation is then however more complex

$$\bar{N}_t = N_i \left(\frac{1 - e^{-Z_i T_i}}{Z_i T_i} \right)$$

- Where T_i is the time between t and t_i (often 1 year)

Stock & catch equation → Russel equation

- Need to have information on weights (direct measurements or from a growth model) to relate back to the original Russel model:

$$Y_t = \sum C_t w_t$$


$$B_t = \sum N_t w_t$$

W_t : Measured directly or obtained from the VBGF

- $B_{t+1} = B_t - Y_t$
 - Note that we ignored totally the recruitment term since only dealt with one cohort. In age/length based stock models we need to make some strong assumptions about the relationship between stock and recruitment


- Used to estimate mortality (Z, F) and abundance (N)
 - Mortality (Z) can be estimated from both age and relative size frequency samples
 - survey or catch data by length
 - length based samples: Need to know growth (K and L_{inf})
 - For fishing mortality (F) we need to know M
 - $F = Z - M$
 - For abundance (N) we also need the total catch removed

Age based measurements



Length based measurements

Snapshots (pseudochorts) most common



- Advantages
 - Populations do have age/size structure
 - Basic biological processes are age/size specific
 - Growth
 - Mortality
 - Fecundity
 - The process of fishing is age/size specific
 - Relatively simple to construct mathematically
 - Model assumption not as strict as in e.g. logistic models
- Disadvantages
 - Sample intensive
 - Data often not available
 - Mostly limited to areas where species diversity is low
 - Have to have knowledge of natural mortality
 - For long term management strategies have to make model assumptions about the relationship between stock and recruitment
 - Often not needed to address the question at hand
- In the tropics do analysis on a pseudocohort (snapshot)